

Design of Plasmonic Catalysts for Efficient H₂ Production from Hydrogen Carrier Molecules

Hiromi YAMASHITA^{1*,2}, Priyanka VERMA¹, Haibo YIN¹, Yasutaka KUWAHARA^{1,2}, Kohsuke MORI^{1,2}

¹Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan.

²Unit of Elements Strategy Initiative for Catalysts & Batteries, Kyoto University, Japan.

E-mail of corresponding author: yamashita@mat.eng.osaka-u.ac.jp

Abstract: The Hydrogen is regarded as a promising fuel. Ammonia borane (NH₃BH₃) and formic acid (HCOOH) containing high hydrogen content are attractive candidates as chemical hydrogen carrier molecules. In the exploitation of highly efficient nanocatalysts, special attention has been drawn to the plasmonic nanostructures, including conventional noble metals and non-conventional semiconductor nanocrystals¹⁻⁸. Plasmonic nanostructures are capable of utilization of abundant solar energy by localized surface plasmon resonance (LSPR). LSPR can be described as the collective oscillation of valence electrons for establishing a resonance between the photons and surface electrons of nanoparticles (NPs), driven by the electromagnetic field of incident light. Such LSPR has promising applications in the energy conversion and storage. In this study, we demonstrated that the plasmonic nanostructures, bimetallic Pd/Ag NPs on SBA-15 mesoporous silica, Pd/MoO_{3-x} nanosheet hybrid, core-shell Au@Pd NPs supported on titanium doped zirconium based amine-functionalized MOFs (UiO-66(Zr_{100-x}Ti_x)), can be used as highly efficient catalysts under visible light for H₂ production from NH₃BH₃ and HCOOH. The moderate control of electron density of Pd metal and surface acid-basic properties of support are key factors to realize efficient H₂ production.



Figure 1. Plasmonic Ag/SBA-15 catalysts.




Figure 2. Plasmonic Pd/MoO_{3-x} catalysts.

Key words: Plasmonic Catalysts, H₂ Production, Hydrogen Carrier, Ag NPs, MOF

References

- [1] Y. Isaka, H. Yamashita, *et al.*, *Angew. Chem. Int. Ed.*, 131 (2019) 5456.
- [2] Y. Kuwahara, H. Yamashita, *et al.*, *J. Am. Chem. Soc.*, 140 (2018) 9203.
- [3] K. Mori, H. Yamashita, *et al.*, *J. Am. Chem. Soc.*, 140 (2018), 8902.
- [4] H. Yamashita, K. Mori, *et al.*, *Chem. Soc. Rev.*, 47 (2018) 8072.
- [5] M. Wen, H. Yamashita, *et al.*, *ACS Energy Lett.*, 2 (2017) 1.
- [6] H. Cheng, H. Yamashita, *et al.*, *J. Am. Chem. Soc.*, 138 (2016) 9316.
- [7] H. Cheng, H. Yamashita, *et al.*, *Angew. Chem. Int. Ed.*, 53 (2014) 2910.
- [8] K. Fuku, H. Yamashita, *et al.*, *Angew. Chem. Int. Ed.*, 52 (2013) 7446.